

Recommended quantity for Zegrow:

Apply Zeogrow any time after draught, stress or natural pests. Overuse will NOT have any adverse effect on the plant.

Application for cultures (per hectare)

- Corn:

1. Application 3 Kg when plant is about 20 cm

2. Application 3 Kg 3-4 weeks later

3. Application 3 Kg before plant is too high to pass over

- Potato:

1. Application 3 Kg when plant has about 15-20 cm, 4 leaves

2. Application 3 Kg 14-21 days later

3. Application 3 Kg after flower

4. Application 2.5 Kg 14-21 days later

Fruit-tree:

- Citrus, Apple, Pears, Plums, Mango, Papaya:

1. Application 3 Kg when first leaves are out

2. Application 3 Kg 14-21 days later

3. Application 3 Kg after flowering

4. Application 3 Kg 21-28 days later

Cereals

- Wheat, Barley, Oats, Triticale etc:

1. Application 3 Kg when plant is about 20 cm

2. Application 3Kg 21 days later

3. Application 3Kg after flowering





- Winter cereal:

- 1. Application 3 Kg in autumn when plant is about 10-15 cm
- 2. Application 3 Kg in spring
- 3. Application 3 Kg 14 days prior to flowering
- 4. Application 3 Kg after flowering

- Raps:

- 1. Application 3 Kg when plant is about 20 cm
- 2. Application 3 Kg 21 days later
- 3. Application 3 Kg after flowering, but before plant is too high to pass over

- Rice:

- 1,2 Application in nursery 2x times (0,3% solution)
- 3. Application 3 Kg immediately after planting in paddy
- 4. Application 3 Kg 14 days later
- 5. Application 3 Kg after flowering

Vegetable:

- Salad:

- 1,2 Application in nursery 2x times (0,3% solution)
- 3. Application 3 Kg immediately after planting in paddy
- 4. Application 3 Kg 7 -14 days later
- 5. Application 3 Kg 7 -14 days later

- Zuchini, Eggplant, Tomato, Paprika, Cucumber etc:

- 1,2 Application in nursery 2x times (0,3% solution)
- 3. Application 3 Kg immediately after planting in paddy
- 4. Application 2 Kg 14-21 days later
- 5. Application 2 Kg 14-21 days later

Repeating with 2 Kg until harvest





- Pumpkin:

1. Application 3 Kg when plant is about 20 cm

2. Application 3 Kg 14-21 days later

3. Application 3 Kg before planting rows close

- Melon:

1. Application 3 Kg when plants are about 20 cm

2. Application 3 Kg 14 days later

3. Application 2 Kg 21 days later

Repeating with 2 Kg until harvest

- Onion, Garlic, Carrot:

1. Application 3 Kg when plant is about 20 cm

2. Application 3 Kg 14-21 days later

3. Application 3 Kg 14-21 days later

Berry:

- Strawberry, Raspberry, Blackberry, etc.

1. Application 3 Kg when plant is about 15 cm high or has its first leaves

2. Application 3 kg 14-21 days later

3. Application 3 kg 14-21 days later

Repeating with 2 Kg until harvest

Soya:

1. Application 3 Kg when plant is about 20 cm

2. Application 3 Kg 14-21 days later

3. Application 3 Kg 14-21 days later, before plant is too high to pass over





Sunflower:

1. Application 3 Kg when plant is about 20 cm

2. Application 3 Kg 14-21 days later

3. Application 3 Kg 14-21 days later, before plant is too high to pass over

Sugar beet:

1. Application 3 Kg when plant is about 20 cm

2. Application 3 Kg 14-21 days later

3. Application 3 Kg 14-21 days later, before plant growing lines are closing

Further treatments: If possible a further treatment 21 days later

Sugar cane:

1. Application 3 Kg after planting or when plant is about 20 cm

2. Application 3 Kg 21-28 days later

3. Application 3 Kg 21-28 days later

Further treatments: if possible without damaging the plant 21-28 days later

Flower:

- Rose, ornamental Flower:

1. Application 3 Kg when first leaves are out

2. Application 3 Kg 14-21 days later

3. Application 3 Kg 14-21 days later, before plant is too high to pass over

Grass:

1. Application 3 Kg after cutting

2. Application 3 Kg 7 -14 days later

3. Application 3 Kg 21 days later





Nursery:

1. Application 3 Kg once there are small sprouts

2. Application 3 Kg 7 days later

3. Application 3 Kg 14-21 days later

Tree, Energy Plant, Miscanthus etc.

1. Application 3 Kg when plant is replanted or leaves are about 20 cm high

2. Application 3 Kg 14-21 days later

3. Application 3 Kg 14-21 days later, before plant is too high to pass over

Application recommendation for other plants ask under:

Info@vmwa.com.au

Volco Minerals PTY LTD

Research&Developement

